



# GIVE YOUR DOG MORE DAYS OF PLAY

Control dog osteoarthritis (OA) pain  
with monthly, long-lasting Librela


**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. Librela should not be used in breeding, pregnant or lactating dogs. Librela should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](http://LibrelaPI.com).

**Librela**<sup>™</sup>  
(bedinvetmab injection)



## What Is Dog OA Pain?

It's a painful, chronic condition that causes your dog to hurt all the time if it's not treated<sup>1</sup>



OA is a type of arthritis that happens when **protective tissue in the joints is worn down**, causing bones to rub against one another.



**Dogs of all ages, sizes, and breeds can have OA pain**—it's not just a condition seen in older dogs.<sup>2</sup>



**OA pain impacts how your dog moves and feels**, and it can make it hard for you both to do the things you love to do together.



OA pain hurts, decreases your dog's willingness to play, and **affects their quality of life**.<sup>1</sup>

**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. Librela should not be used in breeding, pregnant or lactating dogs. Librela should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](http://LibrelaPI.com).





**Librela**<sup>™</sup>  
(bedinvetmab injection)

## Is Your Dog Showing Signs of OA Pain?

An OA diagnosis can be missed in some dogs because the signs may be subtle or overlooked as normal changes related to aging

Nearly 40% of dogs show signs of OA pain, and almost 25% are still undiagnosed<sup>3,4</sup>

Your dog could be suffering from OA pain if they're displaying the following behaviors:



If your dog is showing any of these behaviors, speak to a veterinary professional.



Take the [Dog OA Pain Checklist quiz](#) to see if your dog could be showing signs of OA.

Scan the code with your phone's camera to take the quiz.



**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. Librela should not be used in breeding, pregnant or lactating dogs. Librela should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](http://LibrelaPI.com).

**Librela**<sup>™</sup>  
(bedinvetmab injection)



## Give Your Dog More Days of Play With Librela



**Librela is a first-line treatment option that provides long-term OA pain control** for your dog with a **once-a-month injection** given by your veterinary professional.<sup>5-9</sup>



In clinical studies, Librela was shown to control signs of OA pain in dogs, **which helped them be more active and improved their overall quality of life.**<sup>5,6,10,11</sup>



With long-lasting Librela, **your dog can feel better**, and you can **feel good about their treatment**, so you can **get back to the activities you both love.**<sup>6,10,11</sup>

## Ask Your Vet About Librela



Librela is a monoclonal antibody that specifically targets a key driver of OA pain. It works to reduce pain signals, making it easier for your dog to move and play.<sup>12,13</sup>



Librela reduces OA pain, which can help your dog move and feel better.<sup>6,10,11</sup>



Librela is a once-monthly injection given by a veterinary professional, which means you don't have to worry about giving your dog daily oral medication for their OA pain.



In a clinical study, the most common side effects in dogs taking Librela vs placebo (no medicine) were urinary tract infection, bacterial skin infection, and dermatitis, and were similar for dogs taking placebo.<sup>5</sup>

**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. Librela should not be used in breeding, pregnant or lactating dogs. Librela should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](http://LibrelaPI.com).

**Librela**<sup>™</sup>  
(bedinvetmab injection)



## How to Prepare for Your Vet Visits

Here are some tips to make your appointments with your vet as productive as possible



**Pay attention to the physical and emotional behaviors of your dog before your appointment**

How is your dog acting? Do they have trouble playing, jumping, or climbing stairs? How do they seem to be feeling?



**Take videos and write down notes of your dog's activity and behaviors to share with your vet**

It's important to talk to your vet about how your dog is moving and feeling so they can assess progress.



**Schedule monthly appointments**

Before you leave an appointment, be sure to schedule your next visit to stay on track.



For the best possible results, Librela should be given to your dog by your veterinary professional once a month, every month. Because OA pain is chronic, it needs to be treated continually to see and maintain improvement.

**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. Librela should not be used in breeding, pregnant or lactating dogs. Librela should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](https://www.LibrelaPI.com).

**Librela**<sup>™</sup>  
(bedinvetmab injection)

## See How Librela Can Make a Difference

More than 9 million doses of Librela have been given to dogs in Europe since 2021<sup>14</sup>

Scan the codes with your phone's camera to see real results with Librela.



[Meet Bella](#)



[Meet Emma](#)

### Do you own a cat that could be showing signs of OA?

Ask your veterinarian about Solensia™ (frunevetmab injection), the FIRST and ONLY treatment approved by the FDA for the control of OA pain in cats.

FDA=US Food and Drug Administration.



Scan the code with your phone's camera to learn more.

### IMPORTANT SAFETY INFORMATION FOR

**SOLENSIA:** For use in cats only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Allergic reactions, including anaphylaxis, could potentially occur with self-injection. Solensia should not be used in breeding cats or in pregnant or lactating queens. Solensia should not be administered to cats with a known allergy to frunevetmab. The most common adverse events reported in a clinical study were vomiting and injection site pain. See full Prescribing Information at [SolensiaPI.com](https://www.solensia.com).

**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. Librela should not be used in breeding, pregnant or lactating dogs. Librela should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](https://www.librela.com).




**Librela**<sup>™</sup>  
(bedinvetmab injection)

# Librela Appointment Tracker

Track your dog's progress with Librela.




After your dog receives an injection of Librela, your vet will schedule your next appointment, stamp your dose tracker, and write down any appointment notes or reminders.




DOSE NUMBER	APPT DATE & TIME

NOTES:




DOSE NUMBER	APPT DATE & TIME

NOTES:




DOSE NUMBER	APPT DATE & TIME

NOTES:




DOSE NUMBER	APPT DATE & TIME

NOTES:




DOSE NUMBER	APPT DATE & TIME

NOTES:



DOSE NUMBER	APPT DATE & TIME

NOTES:



DOSE NUMBER	APPT DATE & TIME

NOTES:



# Feel Confident About Choosing Librela to Treat Your Dog's OA Pain



Provides long-term OA pain control for your dog<sup>5,6</sup>



A first-line, once-monthly injection given by your veterinary professional<sup>7-9</sup>



A monoclonal antibody that works to reduce pain signals, making it easier for your dog to move and play<sup>12,13</sup>



Controls signs of OA pain in dogs, which can help them be more active and improve their overall quality of life<sup>5,6,10,11\*</sup>

\*Results from clinical studies.



Get more information at [Librela.com](https://Librela.com).

Scan the code with your phone's camera to explore more.

**References:** **1.** Lascelles BDX et al. *Vet J*. 2019;250:71-78. doi:10.1016/j.tvjl.2019.07.001 **2.** Anderson KL et al. *Front Vet Sci*. 2020;7:200. doi:10.3389/fvets.2020.00220 **3.** Wright A et al. *J Small Anim Pract*. 2022;63(8):609-618. doi:10.1111/jsap.13500 **4.** ZMR: 2023 OA Landscape Wave 6 Analysis: Quantitative Results, Zoetis Inc; June 2023. **5.** Michels GM et al. *Vet Anaesth Analg*. 2023;50(5):446-458. doi:10.1016/j.vaa.2023.06.003 **6.** Corral MJ et al. *Vet Anaesth Analg*. 2021;48:943-955. doi:10.1016/j.vaa.2021.08.001 **7.** Gruen ME et al. *J Am Anim Hosp Assoc*. 2022;58(2):55-76. doi:10.5326/JAAHA-MS-7292 **8.** Monteiro BP et al. *J Small Anim Pract*. 2023;64:177-254. doi:10.1111/jsap.13566 **9.** Cachon T et al. *Front Vet Sci*. 2023;10. doi:10.3389/fvets.2023.1137888 **10.** Brown DC et al. *Am J Vet Res*. 2007;68(6):631-637. doi:10.2460/ajvr.68.6.631 **11.** Brown DC et al. *J Am Vet Med Assoc*. 2008;233(8): 1278-1283. doi:10.2460/javma.233.8.1278 **12.** Keizer RJ et al. *Clin Pharmacokinet*. 2010;49(8):493-507. doi:10.2165/11531280 **13.** Isola Met al. *Vet Comp Orthop Traumatol*. 2011;24(4):279-284. doi:10.3415/VCOT-10-04-0051 **14.** Data on file. Zoetis Inc. November 2023.

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted. © 2024 Zoetis Services LLC. All rights reserved. LIB-00099R1

zoetis

**IMPORTANT SAFETY INFORMATION:** For use in dogs only. Women who are pregnant, trying to conceive or breastfeeding should take extreme care to avoid self-injection. Hypersensitivity reactions, including anaphylaxis, could potentially occur with self-injection. Librela should not be used in breeding, pregnant or lactating dogs. Librela should not be administered to dogs with known hypersensitivity to bedinvetmab. The most common adverse events reported in a clinical study were urinary tract infections, bacterial skin infections and dermatitis. See full Prescribing Information at [LibrelaPI.com](https://LibrelaPI.com).

Librela™  
(bedinvetmab injection)